



A Multiscale Approach to the Asymptotic Behavior of Solutions to Nonlinear Integral Equations with Generalized Heat Kernel

Gastão A. Braga, Jussara M. Moreira,

Departamento de Matemática - Universidade Federal de Minas Gerais

Av. Antônio Carlos, 6627 - Pampulha

31270-901, Belo Horizonte, MG

E-mail: gbraga@mat.ufmg.br, jmoreira@mat.ufmg.br,

Camila F. Souza

Centro Federal de Educação Tecnológica de Minas Gerais

Av. Amazonas, 5.253, Nova Suíça

30421-169, Belo Horizonte, MG

E-mail: camila@des.cefetmg.br

Here we study a class of nonlinear integral equations with a generalized heat kernel, obtaining global existence and uniqueness of the solution, as well as the asymptotic behavior, using the Renormalization Group method. The nonlinearities are classified according to their contribution to the asymptotic behavior. We show that the so called *irrelevant* perturbations do not affect the asymptotic profile of the solution, in the sense that the profile is the same as in the linear case, whereas, by adding *marginal* perturbations, an extra logarithmic factor appears on the decay of the solution.

More specifically, we obtain the asymptotic behavior of solutions to equations of type

$$u(x, t) = \int G(x - y, s(t))f(y)dy + \int_1^t \int G(x - y, s(t) - s(\tau))F(u(y, \tau))dyd\tau, \quad (1)$$

with $x \in \mathbb{R}$ and $t > 1$. By imposing conditions on the kernel without specifying $G = G(x, t)$, we generalize the study of asymptotics for initial value problems. This outlook was adopted in [1, 2] where it is shown that, under certain conditions on G , with $s(t) = t$, the solution $u(x, t)$ to (1) behaves for long time as

$$\frac{A}{t^{1/d}}G\left(\frac{x}{t^{1/d}}, 1\right),$$

where $d > 0$ is such that $G(x, t) = t^{-\frac{1}{d}}G\left(t^{-\frac{1}{d}}x, 1\right)$. We recover and extend the above result using a multiscale approach showing that, if $c(t)$ is a positive function in $L^1_{loc}((1, +\infty))$ of type $t^p + o(t^p)$, with $p > 0$ and

$$s(t) = \int_1^t c(\tau)d\tau = \frac{t^{p+1} - 1}{p + 1} + r(t),$$

then, for $F(u) = \sum_{j \geq \alpha} a_j u^j$ with $\alpha > (p+1+d)/(p+1)$,

$$u(x, t) \sim \frac{A}{t^{(p+1)/d}} G\left(\frac{x}{t^{(p+1)/d}}, \frac{1}{p+1}\right) \text{ when } t \rightarrow \infty.$$

Furthermore, if $F(u) = -\lambda u^\alpha + \sum_{j > \alpha} a_j u^j$ with λ small and positive and $\alpha = (d+p+1)/(p+1)$, then

$$u(x, t) \sim \frac{A}{(t \ln t)^{(p+1)/d}} G\left(\frac{x}{t^{(p+1)/d}}, \frac{1}{p+1}\right) \text{ when } t \rightarrow \infty.$$

Referências

- [1] K. Ishige, T.Kawakami, and K.Kobayashi, *Asymptotics for a Nonlinear Integral Equation with a Generalized Heat Kernel*, “Journal of Evolution Equations”, vol. 14, pp. 749–777, 2014.
- [2] K. Ishige, T.Kawakami, and K.Kobayashi, *Global Solutions for a Nonlinear Integral Equation with a Generalized Heat Kernel*, “Discrete and Continuous Dynamical Systems S”, vol. 7, pp. 767–783, 2014.
- [3] J. Bricmont, A. Kupiainen and G. Lin, *Renormalization Group and Asymptotics of Solutions of Nonlinear Parabolic Equations*, “Comm. Pure Appl. Math.”, 47, 893-922, 1994.
- [4] G. A. Braga, F. Furtado, J. M. Moreira and L. T. Rolla, *Renormalization Group Analysis of Nonlinear Diffusion Equations with Time Dependent Coefficients: Analytical Results*, “Discrete and Continuous Dynamical Systems. Series B”, vol. 7, pp. 699–715, 2007.
- [5] G. A. Braga and J. M. Moreira, *Renormalization Group Analysis of Nonlinear Diffusion Equations with Time Dependent Coefficients and Marginal Perturbations*, “Journal of Statistical Physics”, vol. 148, no. 2, pp. 280–295, 2012.

Keywords: *asymptotic behavior, heat kernel, integral equation, renormalization group*

Agradecimentos: *Agradecemos à CAPES, CNPq e FAPEMIG pelo apoio financeiro.*